



Culturing of *Selenastrum* on diluted composting fluids; conversion of waste to valuable algal biomass in presence of bacteria



Tossavainen Marika ^{a,*}, Nykänen Anne ^a, Valkonen Kalle ^{a,1}, Ojala Anne ^{a,b}, Kostia Silja ^c, Romantschuk Martin ^{a,d}

^a Department of Environmental Sciences, University of Helsinki, Niemenkatu 73, 15140 Lahti, Finland

^b Department of Forest Sciences, P.O. Box 27, 00014 University of Helsinki, Finland

^c Faculty of Technology, Lahti University of Applied Sciences, Ståhlberginkatu 10, 15110 Lahti, Finland

^d Institute of Environmental Sciences, Kazan Federal University, 420008 Kazan, Russia

HIGHLIGHTS

- Composting leachates supported growth of both algae and bacteria in co-cultures.
- Nutrients from composting leachates were efficiently converted to algal biomass.
- Prolonged cultivation time and CO₂ feed enhance the lipid production in co-cultures.
- In commercialization, co-culturing is a realistic alternative.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 6 February 2017

Received in revised form 4 April 2017

Accepted 5 April 2017

Available online 7 April 2017

Keywords:

Microalgae
Wastewater
Fatty acid
Co-culture
Nutrient reduction

ABSTRACT

Growth and fatty acid production of microalga *Selenastrum* sp. with associated bacteria was studied in lab-scale experiments in three composting leachate liquids. Nutrient reduction in cultures was measured at different initial substrate strengths. A small, pilot-scale photobioreactor (PBR) was used to verify lab-scale results. Similar growth conditions supported growth of both *Selenastrum* and bacteria. CO₂ feed enhanced the production of biomass and lipids in PBR (2.4 g L⁻¹ and 17% DW) compared to lab-scale (0.1–1.6 g L⁻¹ and 4.0–6.5% DW) experiments. Also prolonged cultivation time increased lipid content in PBR. At both scales, NH₄-N with an initial concentration of ca. 40 mg L⁻¹ was completely removed from the biowaste leachate. In lab-scale, maximal COD reduction was over 2000 mg L⁻¹, indicating mixotrophic growth of *Selenastrum*. Co-cultures are efficient in composting leachate liquid treatment, and conversion of waste to biomass is a promising approach to improve the bioeconomy of composting plants.

© 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Microalgae are promising as beneficial organisms in production of renewable energy or valuable metabolites such as fatty acids

* Corresponding author.

E-mail address: marika.tossavainen@helsinki.fi (M. Tossavainen).

¹ Present address: Kyrö Distillery Company, Rye Rye Oy, Oltermannintie 6, 61500 Isokyrö, Finland.